

Diagnoses of HIV and/or AIDS Reported in Hamilton County

Diagnoses of HIV infection as of September 30, 2012, by disease status at initial diagnosis and by selected characteristics, Hamilton County

Characteristic	2011 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	27.0	104	73%	76	72%	20	74%	8	89%
Females	9.1	38	27%	30	28%	7	26%	1	11%
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	20.8	12	8%	11	10%	1	4%	-	-
20-24	55.4	33	23%	26	25%	7	26%	-	-
25-29	37.9	22	15%	20	19%	2	7%	-	-
30-34	31.9	16	11%	9	8%	6	22%	1	11%
35-39	23.2	11	8%	6	6%	3	11%	2	22%
40-44	25.7	13	9%	11	10%	-	-	2	22%
45-49	27.2	16	11%	11	10%	3	11%	2	22%
50-54	18.0	11	8%	8	8%	3	11%	-	-
55-64	6.3	6	4%	3	3%	2	7%	1	11%
65+	*	2	1%	1	1%	-	-	1	11%
Race/Ethnicity									
White, not Hispanic	8.3	45	32%	34	32%	5	19%	6	67%
Black/African American, not Hispanic	44.8	92	65%	68	64%	21	78%	3	33%
Hispanic/Latino	*	3	2%	2	2%	1	4%	-	-
Asian/Pacific Islander	*	2	1%	2	2%	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Race/Sex									
White, not Hispanic Males	14.0	37	26%	26	25%	5	19%	6	67%
White, not Hispanic Females	2.9	8	6%	8	8%	-	-	-	-
Black/African American, not Hispanic Males	68.2	64	45%	48	45%	14	52%	2	22%
Black/African American, not Hispanic Females	25.1	28	20%	20	19%	7	26%	1	11%
Hispanic/Latino Males	*	2	1%	1	1%	1	4%	-	-
Hispanic/Latina Females	*	1	1%	1	1%	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-	-	-
Asian/Pacific Islander Females	*	1	1%	1	1%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Total	17.7	142		106		27		9	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate and/or percent not calculated due to small numbers.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Diagnoses of HIV infection as of September 30, 2012, by disease status at initial diagnosis and by selected characteristics, Hamilton County

Transmission Category	2011 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
	HIV infection		HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	60	58%	45	59%	12	60%	3	38%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	3	3%	2	3%	-	-	1	13%
Heterosexual contact	3	3%	3	4%	-	-	-	-
Other/unknown	38	37%	26	34%	8	40%	4	50%
Subtotal	104	100%	76	100%	20	100%	8	100%
Female adult or adolescent								
Injection drug use	1	3%	1	3%	-	-	-	-
Heterosexual contact	10	26%	9	30%	1	14%	-	-
Other/unknown	27	71%	20	67%	6	86%	1	100%
Subtotal	38	100%	30	100%	7	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	142		106		27		9	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

**Diagnosis of HIV infection as of September 30, 2012, by race/ethnicity and transmission category,
Hamilton County**

Transmission Category	2011 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	22	59%	35	55%	2	100%	1	100%	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	5%	1	2%	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	3	5%	-	-	-	-	-	-	-	-
Other/unknown	13	35%	25	39%	-	-	-	-	-	-	-	-
Subtotal	37	100%	64	100%	2	100%	1	100%	-	-	-	-
Female adult or adolescent												
Injection drug use	1	13%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	38%	6	21%	-	-	1	100%	-	-	-	-
Other/unknown	4	50%	22	79%	1	100%	-	-	-	-	-	-
Subtotal	8	100%	28	100%	1	100%	1	100%	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	45		92		3		2		-		-	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported diagnoses of HIV infection as of September 30, 2012, by year of diagnosis and cumulative diagnoses by selected characteristics, Hamilton County

Diagnosis of HIV Infection by Year

Characteristic	2007			2008			2009			2010		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex												
Males	25.2	97	82%	23.9	92	75%	30.4	117	73%	30.4	117	76%
Females	5.0	21	18%	7.2	30	25%	10.3	43	27%	8.6	36	24%
Age at diagnosis (yr)												
<13	*	1	1%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	13.9	8	7%	12.1	7	6%	27.7	16	10%	31.2	18	12%
20-24	36.9	22	19%	57.0	34	28%	62.1	37	23%	57.0	34	22%
25-29	25.9	15	13%	19.0	11	9%	37.9	22	14%	46.6	27	18%
30-34	31.9	16	14%	18.0	9	7%	35.9	18	11%	33.9	17	11%
35-39	23.2	11	9%	29.6	14	11%	21.1	10	6%	29.6	14	9%
40-44	33.7	17	14%	49.5	25	20%	23.8	12	8%	29.7	15	10%
45-49	23.8	14	12%	13.6	8	7%	27.2	16	10%	32.3	19	12%
50-54	11.5	7	6%	16.4	10	8%	24.6	15	9%	13.1	8	5%
55-64	6.3	6	5%	*	4	3%	13.6	13	8%	*	1	1%
65+	*	1	1%	*	-	-	*	1	1%	*	-	-
Race/Ethnicity												
White, not Hispanic	6.8	37	31%	6.1	33	27%	7.0	38	24%	8.3	45	29%
Black/African American, not Hispanic	36.5	75	64%	40.4	83	68%	56.0	115	72%	50.6	104	68%
Hispanic/Latino	*	4	3%	24.3	5	4%	*	4	3%	*	-	-
Asian/Pacific Islander	*	1	1%	*	-	-	*	1	1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	1	1%	*	2	1%	*	4	3%
Race/Sex												
White, not Hispanic Males	12.1	32	27%	9.8	26	21%	11.3	30	19%	13.2	35	23%
White, not Hispanic Females	1.8	5	4%	2.5	7	6%	2.9	8	5%	3.6	10	7%
Black/African American, not Hispanic Males	66.0	62	53%	63.9	60	49%	86.3	81	51%	85.2	80	52%
Black/African American, not Hispanic Females	11.7	13	11%	20.6	23	19%	30.5	34	21%	21.5	24	16%
Hispanic/Latino Males	*	2	2%	45.7	5	4%	*	3	2%	*	-	-
Hispanic/Latina Females	*	2	2%	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	1	1%	*	2	1%	*	4	3%
Total	14.7	118		15.2	122		19.9	160		19.1	153	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported cases of HIV infection diagnosed since the beginning of the epidemic through 2011.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate and/or percent not calculated due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported diagnoses of HIV infection as of September 30, 2012, by year of diagnosis and cumulative diagnoses by transmission category, Hamilton County

Diagnosis of HIV Infection by Year

Transmission Category	2007		2008		2009		2010	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	71	73%	62	67%	81	69%	83	71%
Injection drug use (IDU)	1	1%	3	3%	2	2%	-	-
Male-to-male sexual contact and IDU	2	2%	3	3%	2	2%	3	3%
Heterosexual contact	15	15%	7	8%	6	5%	10	9%
Other/unknown	8	8%	17	18%	26	22%	21	18%
Subtotal	97	100%	92	100%	117	100%	117	100%
Female adult or adolescent								
Injection drug use	1	5%	4	13%	7	16%	3	8%
Heterosexual contact	15	75%	15	50%	13	30%	11	31%
Other/unknown	4	20%	11	37%	23	53%	22	61%
Subtotal	20	100%	30	100%	43	100%	36	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-
Total	118		122		160		153	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported cases of HIV infection diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported HIV (not AIDS) diagnoses as of September 30, 2012, by year of HIV diagnosis and by selected characteristics, Hamilton County

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	62	82%	52	74%	62	71%	68	73%	76	72%
Females	14	18%	18	26%	25	29%	25	27%	30	28%
Age at HIV diagnosis (yr)										
<13	1	1%	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	6	8%	3	4%	11	13%	17	18%	11	10%
20-24	13	17%	23	33%	26	30%	28	30%	26	25%
25-29	11	14%	7	10%	11	13%	13	14%	20	19%
30-34	13	17%	5	7%	11	13%	9	10%	9	8%
35-39	8	11%	7	10%	3	3%	7	8%	6	6%
40-44	9	12%	15	21%	4	5%	6	6%	11	10%
45-49	7	9%	5	7%	8	9%	9	10%	11	10%
50-54	6	8%	3	4%	7	8%	4	4%	8	8%
55-64	2	3%	2	3%	6	7%	-	-	3	3%
65+	-	-	-	-	-	-	-	-	1	1%
Race/Ethnicity										
White, not Hispanic	26	34%	16	23%	21	24%	29	31%	34	32%
Black/African American, not Hispanic	47	62%	50	71%	62	71%	60	65%	68	64%
Hispanic/Latino	2	3%	4	6%	3	3%	-	-	2	2%
Asian/Pacific Islander	1	1%	-	-	-	-	-	-	2	2%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	1%	4	4%	-	-
Race/Sex										
White, not Hispanic Males	22	29%	12	17%	16	18%	21	23%	26	25%
White, not Hispanic Females	4	5%	4	6%	5	6%	8	9%	8	8%
Black/African American, not Hispanic Males	38	50%	36	51%	42	48%	45	48%	48	45%
Black/African American, not Hispanic Females	9	12%	14	20%	20	23%	15	16%	20	19%
Hispanic/Latino Males	1	1%	4	6%	3	3%	-	-	1	1%
Hispanic/Latina Females	1	1%	-	-	-	-	-	-	1	1%
Asian/Pacific Islander Males	1	1%	-	-	-	-	-	-	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	1%	4	4%	-	-
Total	76		70		87		93		106	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported HIV (not AIDS) diagnoses as of September 30, 2012, by year of HIV diagnosis and by transmission category, Hamilton County

Transmission Category	Year of HIV (not AIDS) Diagnosis									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	43	69%	31	60%	47	76%	48	71%	45	59%
Injection drug use (IDU)	-	-	3	6%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	3%	-	-	2	3%	2	3%	2	3%
Heterosexual contact	9	15%	6	12%	-	-	7	10%	3	4%
Other/unknown	8	13%	12	23%	13	21%	11	16%	26	34%
Subtotal	62	100%	52	100%	62	100%	68	100%	76	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	3	12%	3	12%	1	3%
Heterosexual contact	11	85%	8	44%	6	24%	8	32%	9	30%
Other/unknown	2	15%	10	56%	16	64%	14	56%	20	67%
Subtotal	13	100%	18	100%	25	100%	25	100%	30	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-
Total	76		70		87		93		106	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported cases of AIDS as of September 30, 2012, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Hamilton County

Characteristic	Year of AIDS Diagnosis									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	40	83%	64	77%	74	72%	70	81%	56	76%
Females	8	17%	19	23%	29	28%	16	19%	18	24%
Age at AIDS diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	4	5%	6	6%	-	-	2	3%
20-24	6	13%	8	10%	11	11%	7	8%	15	20%
25-29	4	8%	5	6%	16	16%	21	24%	7	9%
30-34	5	10%	6	7%	10	10%	12	14%	13	18%
35-39	7	15%	15	18%	13	13%	9	10%	8	11%
40-44	9	19%	22	27%	20	19%	10	12%	3	4%
45-49	9	19%	10	12%	10	10%	12	14%	8	11%
50-54	2	4%	9	11%	11	11%	8	9%	10	14%
55-64	5	10%	4	5%	5	5%	6	7%	7	9%
65+	1	2%	-	-	1	1%	1	1%	1	1%
Race/Ethnicity										
White, not Hispanic	13	27%	26	31%	29	28%	25	29%	22	30%
Black/African American, not Hispanic	33	69%	55	66%	67	65%	60	70%	51	69%
Hispanic/Latino	2	4%	1	1%	3	3%	-	-	1	1%
Asian/Pacific Islander	-	-	-	-	1	1%	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	1%	3	3%	1	1%	-	-
Race/Sex										
White, not Hispanic Males	11	23%	24	29%	25	24%	23	27%	18	24%
White, not Hispanic Females	2	4%	2	2%	4	4%	2	2%	4	5%
Black/African American, not Hispanic Males	28	58%	38	46%	46	45%	46	53%	37	50%
Black/African American, not Hispanic Females	5	10%	17	20%	21	20%	14	16%	14	19%
Hispanic/Latino Males	1	2%	1	1%	1	1%	-	-	1	1%
Hispanic/Latina Females	1	2%	-	-	2	2%	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	1	1%	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	1%	3	3%	1	1%	-	-
Total	48		83		103		86		74	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported cases of AIDS as of September 30, 2012, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Hamilton County

Transmission Category	Year of AIDS Diagnosis									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	29	73%	46	72%	48	65%	52	74%	34	61%
Injection drug use (IDU)	1	3%	1	2%	2	3%	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	1	2%	3	4%	3	4%	2	4%
Heterosexual contact	7	18%	7	11%	9	12%	3	4%	1	2%
Other/unknown	2	5%	9	14%	12	16%	12	17%	19	34%
Subtotal	40	100%	64	100%	74	100%	70	100%	56	100%
Female adult or adolescent										
Injection drug use	-	-	1	5%	5	17%	1	6%	2	12%
Heterosexual contact	6	75%	16	84%	14	48%	4	25%	5	29%
Other/unknown	2	25%	2	11%	10	34%	11	69%	10	59%
Subtotal	8	100%	19	100%	29	100%	16	100%	17	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	1	100%
Total	48		83		103		86		74	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Persons Living with a Diagnosis of HIV and/or AIDS Reported in Hamilton County

Reported persons living with a diagnosis of HIV infection as of September 30, 2012, by current disease status and by selected characteristics, Hamilton County

Characteristic	Living with a diagnosis of HIV infection			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	476.3	1,835	79%	898	77%	937	81%
Females	115.3	481	21%	267	23%	214	19%
Age at diagnosis (yr)							
<13	*	2	<1%	1	<1%	1	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	43.3	25	1%	23	2%	2	<1%
20-24	218.1	130	6%	97	8%	33	3%
25-29	296.6	172	7%	117	10%	55	5%
30-34	339.2	170	7%	97	8%	73	6%
35-39	513.4	243	10%	139	12%	104	9%
40-44	576.1	291	13%	134	12%	157	14%
45-49	757.7	446	19%	193	17%	253	22%
50-54	601.3	367	16%	184	16%	183	16%
55-64	395.0	379	16%	142	12%	237	21%
65+	84.2	90	4%	38	3%	52	5%
Race/Ethnicity							
White, not Hispanic	167.8	912	39%	465	40%	447	39%
Black/African American, not Hispanic	647.3	1,330	57%	658	56%	672	58%
Hispanic/Latino	179.6	37	2%	20	2%	17	1%
Asian/Pacific Islander	65.3	11	<1%	7	1%	4	<1%
American Indian/Alaska Native	*	2	<1%	-	-	2	<1%
Unknown	*	24	1%	15	1%	9	1%
Race/Sex							
White, not Hispanic Males	298.4	789	34%	383	33%	406	35%
White, not Hispanic Females	44.1	123	5%	82	7%	41	4%
Black/African American, not Hispanic Males	1,052.1	988	43%	482	41%	506	44%
Black/African American, not Hispanic Females	306.6	342	15%	176	15%	166	14%
Hispanic/Latino Males	255.8	28	1%	15	1%	13	1%
Hispanic/Latina Females	93.1	9	<1%	5	<1%	4	<1%
Asian/Pacific Islander Males	108.5	9	<1%	6	1%	3	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	-	-	2	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	24	1%	15	1%	9	1%
Total	288.6	2,316		1,165		1,151	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate and/or percent not calculated due to small numbers.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported persons living with a diagnosis of HIV infection as of September 30, 2012, by current disease status and by transmission category, Hamilton County

Transmission Category	Living with a diagnosis of HIV infection		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,231	67%	567	63%	664	71%
Injection drug use (IDU)	58	3%	26	3%	32	3%
Male-to-male sexual contact and IDU	75	4%	27	3%	48	5%
Heterosexual contact	109	6%	46	5%	63	7%
Other/unknown	354	19%	227	25%	127	14%
Subtotal	1,827	100%	893	100%	934	100%
Female adult or adolescent						
Injection drug use	44	9%	18	7%	26	12%
Heterosexual contact	252	53%	129	49%	123	58%
Other/unknown	178	38%	116	44%	62	29%
Subtotal	474	100%	263	100%	211	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	67%	7	78%	3	50%
Other/unknown	5	33%	2	22%	3	50%
Subtotal	15	100%	9	100%	6	100%
Total	2,316		1,165		1,151	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported persons living with a diagnosis of HIV infection as of September 30, 2012, by year and by transmission category, Hamilton County

Transmission Category	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	611	78%	586	60%	14	50%	7	78%	1	50%	12	63%
Injection drug use (IDU)	11	1%	42	4%	5	18%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	35	4%	39	4%	-	-	-	-	1	50%	-	-
Heterosexual contact	10	1%	94	10%	4	14%	1	11%	-	-	-	-
Other/unknown	120	15%	221	23%	5	18%	1	11%	-	-	7	37%
Subtotal	787	100%	982	100%	28	100%	9	100%	2	100%	19	100%
Female adult or adolescent												
Injection drug use	19	15%	24	7%	-	-	-	-	-	-	1	20%
Heterosexual contact	62	50%	179	53%	7	78%	2	100%	-	-	2	40%
Other/unknown	42	34%	132	39%	2	22%	-	-	-	-	2	40%
Subtotal	123	100%	335	100%	9	100%	2	100%	-	-	5	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	10	77%	-	-	-	-	-	-	-	-
Other/unknown	2	100%	3	23%	-	-	-	-	-	-	-	-
Subtotal	2	100%	13	100%	-	-	-	-	-	-	-	-
Total	912		1,330		37		11		2		24	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported persons living with a diagnosis of HIV infection as of September 30, 2012, by year and by selected characteristics, Hamilton County

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	1,422	81%	1,506	80%	1,618	80%	1,733	80%	1,835	79%
Females	343	19%	371	20%	410	20%	444	20%	481	21%
Age at diagnosis (yr)										
<13	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	4	<1%	4	<1%	5	<1%	14	1%	25	1%
20-24	21	1%	35	2%	67	3%	99	5%	130	6%
25-29	67	4%	96	5%	125	6%	151	7%	172	7%
30-34	109	6%	119	6%	136	7%	156	7%	170	7%
35-39	188	11%	197	10%	212	10%	227	10%	243	10%
40-44	244	14%	255	14%	265	13%	279	13%	291	13%
45-49	385	22%	405	22%	420	21%	436	20%	446	19%
50-54	318	18%	328	17%	341	17%	353	16%	367	16%
55-64	340	19%	349	19%	367	18%	372	17%	379	16%
65+	86	5%	86	5%	87	4%	87	4%	90	4%
Race/Ethnicity										
White, not Hispanic	757	43%	790	42%	826	41%	869	40%	912	39%
Black/African American, not Hispanic	954	54%	1,028	55%	1,136	56%	1,239	57%	1,330	57%
Hispanic/Latino	26	1%	30	2%	34	2%	34	2%	37	2%
Asian/Pacific Islander	8	<1%	8	<1%	9	<1%	9	<1%	11	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	18	1%	19	1%	21	1%	24	1%	24	1%
Race/Sex										
White, not Hispanic Males	665	38%	691	37%	720	36%	754	35%	789	34%
White, not Hispanic Females	92	5%	99	5%	106	5%	115	5%	123	5%
Black/African American, not Hispanic Males	715	41%	768	41%	845	42%	924	42%	988	43%
Black/African American, not Hispanic Females	239	14%	260	14%	291	14%	315	14%	342	15%
Hispanic/Latino Males	19	1%	23	1%	26	1%	26	1%	28	1%
Hispanic/Latina Females	7	<1%	7	<1%	8	<1%	8	<1%	9	<1%
Asian/Pacific Islander Males	7	<1%	7	<1%	8	<1%	8	<1%	9	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	18	1%	19	1%	21	1%	24	1%	24	1%
Total	1,765		1,877		2,028		2,177		2,316	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Reported persons living with a diagnosis of HIV infection as of September 30, 2012, by year and by transmission category, Hamilton County

Living with a Diagnosis of HIV Infection

Transmission Category	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	954	67%	1,013	68%	1,091	68%	1,172	68%	1,231	67%
Injection drug use (IDU)	53	4%	56	4%	58	4%	58	3%	58	3%
Male-to-male sexual contact and IDU	65	5%	68	5%	70	4%	73	4%	75	4%
Heterosexual contact	83	6%	90	6%	96	6%	106	6%	109	6%
Other/unknown	259	18%	271	18%	295	18%	316	18%	354	19%
Subtotal	1,414	100%	1,498	100%	1,610	100%	1,725	100%	1,827	100%
Female adult or adolescent										
Injection drug use	30	9%	34	9%	40	10%	43	10%	44	9%
Heterosexual contact	205	61%	219	60%	231	57%	242	55%	252	53%
Other/unknown	101	30%	111	30%	132	33%	152	35%	178	38%
Subtotal	336	100%	364	100%	403	100%	437	100%	474	100%
Child (<13 yrs at diagnosis)										
Perinatal	10	67%	10	67%	10	67%	10	67%	10	67%
Other/unknown	5	33%	5	33%	5	33%	5	33%	5	33%
Subtotal	15	100%	15	100%	15	100%	15	100%	15	100%
Total	1,765		1,877		2,028		2,177		2,316	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.